

<b>Patient Name</b> SAMPLEREPORT,FPSAU NORMAL	<b>Patient ID</b> SA00049266	<b>Age</b> 56	<b>Gender</b> M	<b>Order #</b> SA00049266
<b>Ordering Phys</b>				<b>DOB</b> 05/16/1956
<b>Client Order #</b> SA00049266	<b>Account Information</b>			<b>Report Notes</b>
<b>Collected</b> 09/25/2012 10:49	C7028846-DLMP ROCHESTER 3050 SUPERIOR DRIVE ROCHESTER,MN 55901			
<b>Printed</b> 09/25/2012 14:42				

Test	Flag	Results	Unit	Reference Value	Perform Site*
PSA, Ultrasensitive		0.251	ng/mL	0.000 - 4.000	Y06 0

Roche ECLIA methodology.

According to the American Urological Association, Serum PSA should decrease and remain at undetectable levels after radical prostatectomy. The AUA defines biochemical recurrence as an initial PSA value 0.200 ng/mL or greater followed by a subsequent confirmatory PSA value of 0.200 ng/mL or greater. Values obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

\* Performing Site:

Y060	LabCorp Burlington 1447 York Court Burlington, NC 27215-2230	Lab Director:
------	-----------------------------------------------------------------	---------------

<b>Patient Name</b> SAMPLEREPORT,FPSAU NORMAL	<b>Collection Date and Time</b> 09/25/2012 10:49	<b>Report Status</b> Final
Page 1 of 1		** End of Report **

\* Report times for Mayo performed tests are CST/CDT