MAYO CLINIC
Mayo Medical Laboratories

Laboratory Service Report 1-800-533-1710

Patient Name	Patient ID	Age	Gender	Order #
SAMPLEREPORT,HTG1	SA00054570	46	F	SA00054570
Ordering Phys CLIENT,CLIENT				**DOB** 06/10/1966
Client Order # SA00054570	**Account Information**			**Report Notes**
Collected 03/07/2013	C7028846-DLMP Rochester 3050 Superior Drive Rochester, MN 55901			
Printed 03/26/2013 14:46				

Test	Flag	Results	Unit	Reference Value	Perform Site*
Thyroglobulin, Tumor Marker, S			REPORTED 03/07/2013 10:02		
Thyroglobulin, Tumor Marker, S	H	2001	ng/mL		SDL

Quantitation of thyroglobulin may be unreliable due to the presence of anti-thyroglobulin antibodies.
-- REFERENCE VALUE --
<=33
Athyrotic individuals normally have hTg values <=2.

Thyroglobulin Antibody Screen, S	H	22	IU/mL	<22	SDL

Thyroglobulin concentrations >2000 ng/mL may lead to falsely elevated thyroglobulin antibody concentrations and should be interpreted with caution.
The thyroglobulin testing method is an immunoenzymatic assay manufactured by Beckman Coulter Inc. and performed on the Unicel DXI 800.
The thyroglobulin antibody testing method is an electrochemiluminescence assay manufactured by Roche Diagnostics Inc. and performed on the Modular or Cobas System.

Values obtained from different assay methods or kits may be different and cannot be used interchangeably.

The results cannot be interpreted as absolute evidence for the presence or absence of malignant disease.

Specimens with thyroglobulin concentrations greater than 250,000 ng/mL may give falsely lower results.

* Performing Site:

SDL	Mayo Clinic Laboratories - Rochester Superior Drive 3050 Superior Dr. NW Rochester, MN 55901	Lab Director: Franklin R. Cockerill, III, M.D.

* Report times for Mayo performed tests are CST/CDT