



Plasminogen Activator Inhibitor Antigen,  
Plasma

Patient ID <b>SA00062835</b>	Patient Name <b>TESTINGRNV, RUO PAI1</b>	Birth Date <b>1967-10-21</b>	Gender <b>F</b>	Age <b>45</b>
Order Number <b>SA00062835</b>	Client Order Number <b>SA00062835</b>	Ordering Physician <b>Client, Client</b>	Report Notes	
Account Information <b>C7028846 DLMP Rochester</b>		Collected <b>29 Sep 2013 13:56</b>		

**PAI-1 Ag, P**

**MCR**



**<3 ng/mL**

**Reference Value**  
3-72

**Low**

Decreased plasma PAI-1 antigen may occur with the intravascular coagulation and fibrinolysis (ICF/DIC), or may be observed in up to 6.8% of apparently healthy adults. Persistently undetectable PAI-1 antigen rarely may be associated with a congenital bleeding disorder. Suggest clinical correlation and if indicated, repeat testing with careful attention to an early morning, fasting plasma sample.

**ADDITIONAL INFORMATION**

The plasma PAI-1 antigen normal range is based on fasting morning samples in adults. PAI-1 levels follow a diurnal variation, with peak levels occurring in the early morning and nadir levels in the afternoon.

**Received:** 30 Sep 2013 14:05

**Reported:** 30 Sep 2013 14:20

**Performing Site Legend**

Code	Laboratory	Address
MCR	Mayo Clinic Dept. of Lab Med and Pathology	200 First Street SW, Rochester, MN 55905