

Lymphocyte Proliferation Panel for Mitogens and Antigens

Patient ID SA00057307	Patient Name SAMPLEREPOR, LPMAF	Birth Date 1966-06-10	Gender F	Age 46
Order Number SA00057307	Client Order Number SA00057307	Ordering Physician Client, Client	Report Notes	
Account Information C7028846 DLMP Rochester		Collected 08 May 2013 00:00		

Lymphocyte Proliferation Panel

Lymphocyte Proliferation, Mitogens

Interpretation

1 MCR

Normal and robust lymphocyte proliferative responses to PHA and PWM.

ADDITIONAL INFORMATION

Data are expressed as % proliferating cells of total specific cell population. The % Day 0 viability of the sample was determined using a flow cytometry assay which includes individual assessment of viable, apoptotic and dead cells. This method differs from the commonly used method of trypan blue dye exclusion which only identifies dead cells, and counts apoptotic cells along with the viable cells, resulting in an apparent higher cell viability. However, apoptotic cells do not contribute to cell proliferation and therefore accurate measurement of only viable cells provides meaningful information on the cells involved in stimulation and proliferative response. Strongly recommend using "critical ambient shipping boxes" available through Mayo Medical Laboratories (MML) inventory to ensure optimal transport of critical samples used for functional cellular assays.

Viab of Lymphs at Day 0

MCR

79.4 %

 Reference Value
 ≥75.0

Max Prolif of PWM as % CD45

MCR

8.7 %

 Reference Value
 ≥4.5

Max Prolif of PWM as % CD3

MCR

10.2 %

 Reference Value
 ≥3.5

Max Prolif of PWM as % CD19

MCR

9.4 %

 Reference Value
 ≥3.9

Max Prolif of PHA as % CD45

MCR

58.3 %

 Reference Value
 ≥49.9

Max Prolif of PHA as % CD3

MCR

69.9 %

 Reference Value
 ≥58.5

Mitogen Comment

MCR

Lymphocyte proliferative responses are affected by sample age. Samples received between 24–48 hours post-collection can show significant decrease in lymphocyte proliferative responses. Caution should be used when interpreting the results and clinical correlation is strongly recommended. Suggest repeat testing when clinically appropriate.

Lymphocyte Proliferation, Antigens

Interpretation

1 MCR

Normal and robust proliferative responses to antigens - Candida (CA) and Tetanus toxoid (TT).

ADDITIONAL INFORMATION

Data are expressed as % proliferating cells of total specific cell population. The % Day 0 viability of the sample was determined using a flow cytometry assay which includes individual assessment of viable, apoptotic and dead cells. This method

differs from the commonly used method of trypan blue dye exclusion which only identifies dead cells, and counts apoptotic cells along with the viable cells, resulting in an apparent higher cell viability. However, apoptotic cells do not contribute to cell proliferation and therefore accurate measurement of only viable cells provides meaningful information on the cells involved in stimulation and proliferative response. Strongly recommend using "critical ambient shipping boxes" available through Mayo Medical Laboratories (MML) inventory to ensure optimal transport of critical samples used for functional cellular assays.

Performing Site Legend

Code	Laboratory	Address
MCR	Mayo Clinic Dept. of Lab Med and Pathology	200 First Street SW, Rochester, MN 55905



Patient ID SA00057307	Patient Name SAMPLEREPORT, LPMAF	Birth Date 1966-06-10	Gender F	Age 46
Order Number SA00057307	Client Order Number SA00057307	Ordering Physician Client, Client	Report Notes	
Account Information C7028846 DLMP Rochester		Collected 08 May 2013 00:00		

Viab of Lymphs at Day 0

79.4 %

MCR
Reference Value
≥75.0

Max Prolif of TT as % CD3

18.7 %

MCR
Reference Value
≥3.3

Max Prolif of CA as % CD45

12.7 %

MCR
Reference Value
≥5.7

Antigen Comment

Lymphocyte proliferative responses are affected by sample age. Samples received between 24–48 hours post-collection can show significant decrease in lymphocyte proliferative responses. Caution should be used when interpreting the results and clinical correlation is strongly recommended. Suggest repeat testing when clinically appropriate.

MCR

Max Prolif of CA as % CD3

11.8 %

MCR
Reference Value
≥3.0

Received: 09 May 2013 15:22

Reported: 13 Jun 2013 12:42

Max Prolif of TT as % CD45

18.3 %

MCR
Reference Value
≥5.2

Laboratory Notes

- 1 Analyte Specific Reagent: This test was developed and its performance characteristics determined by Mayo Clinic. It has not been cleared or approved by the U.S. Food and Drug Administration.

Performing Site Legend

Code	Laboratory	Address
MCR	Mayo Clinic Dept. of Lab Med and Pathology	200 First Street SW, Rochester, MN 55905